



Guidance for children returning to school or day care - during COVID-19 pandemic

COVID-19 will be part of our lives for the long term and measures to balance risk of infection and risk of continued confinement will need to be evaluated continually.

Lifting of restrictive public health measures including gradual reopening of schools will occur – at different times and with varying interventions. The risk of acquiring COVID-19 varies across Canada and in each jurisdiction. It is a dynamic process and it will be important to continue to follow public health recommendations over time. The return to school in Quebec, planned for May 11 2020 is currently optional. In Ontario, schools remain officially closed until May 31st. This decision will be re-evaluated in the coming weeks and school closure may continue.

Goal:

- Mimimize risk of serious illness and death in infants, children and adolescents related to COVID 19
- Identify the children at greater risk of serious illness and death from COVID-19

What is known about COVID-19 in children:

Current epidemiologic data and studies consistently report that infants, children and adolescents do not develop serious disease from COVID 19 compared with adults and elderly. Even in studies looking at the impact of pediatric underlying co-morbidities on COVID, the disease remained mild to moderate. Death from COVID 19 in pediatrics has been extremely rare. A scan of the literature up to April 26 reveals that proportions of children infected with COVID-19 range between 1.2% to 5% (CPS statement, see ref).

The vast majority of children infected with COVID-19 have had mild to moderate disease, with clinical symptoms very similar to any other viral upper respiratory tract infection.

In addition, current data suggests that school transmission may not be common. In a recent study (NWS report), where there were 18 confirmed cases, no teacher or staff member contracted COVID-19 from any of the initial school cases (735 students and 128 staff were close contacts of these initial 18 cases).

As of May 2020, we are currently in a period where there is no vaccination, herd immunity or readily available sero-diagnostic testing. The return to school or lifting of restrictive public health measures should be associated with the continued promotion of principles of infection prevention and control within the schools and outside the home. Physical distancing, hand hygiene, cloth masking, cough etiquette and limiting large gatherings should be maintained as best as possible during this time.

Specific populations:

At this time, it may be cautious to consider certain specific groups of patients as potentially being more vulnerable, although not reported, to developing serious illness with COVID-19. These may include the following patients:

High level immunosuppression from medication:





- High dose corticosteroids prednisone equivalent > 20 mg per day or 2 mg/kg/day for > 14 days
- Receiving certain monoclonal antibody biologics (ex: Anti-TNF, rituximab, Xolair)
- Patients receiving combination immunosuppressive therapies (> 1 class agents)

Higher risk of serious illness due to underlying condition:

- Malignancy on active chemotherapy
- Post HSCT (especially in first 2 year and on immunosuppression and/or evidence of GVHD)
- Post Solid organ transplant (within 2 months)
- Combined primary immunodeficiency (ex: SCID, CID, etc)
- Certain patients with chronic lung disease
 - On > 0.5-1 mg/kg/day for > 2 weeks and/or pulse IV corticosteroids
 - o Receiving supplemental O2 and/or BIPAP, tracheostomy
 - o FEV1 < 80%, or significant pulmonary decline in past year
 - Severe asthma (admitted to PICU in past 24 months, frequent ED visits) or poorly controlled asthma
 - o Bulbar dysfunction or impaired airway clearance
 - Cystic fibrosis
- Severe chronic liver disease
- HIV with CD4 T cells < 200 or < 15% in children < 5 yo
- Certain patient with cardiac conditions:
 - unrepaired cyanotic congenital heart disease, single ventricle palliation, severe pulmonary hypertension, post-heart transplant, moderately to severely decreased heart function/failure, severe hypertrophic cardiomyopathy, unstable ventricular arrhythmias, Kawasaki disease with giant aneurysms or severe coronary stenosis

For consideration

- Patients with cystic fibrosis
- < 2 yo with history of prematurity and a diagnosis of bronchopulmonary dysplasia (especially if on home oxygen therapy or PPNH)
- Neurological or neurodevelopmental conditions (neuromuscular, neurovascular, neurodegenerative and neurodevelopmental conditions and seizure disorders).
- Patients with sickle cell anemia

Importance of optimizing underlying medical condition:

SARS-CoV2, like any other viral infection can worsen an underlying disease or trigger a flare of an underlying condition. As such, it is <u>crucial</u> at this time that children with chronic medical conditions continue their maintenance medication and optimize the underlying illness.

For example, it has <u>not</u> been shown that Type 1 diabetes or asthma patients have more critical COVID-19 disease. However, a viral infection could lead to acute diabetes related complications during illness or an acute asthma exacerbation.

For consideration:

 Delay school start for those patients until school has organized physical distancing/IPAC measures as to minimize risk.





- If your child is healthy but living with a vulnerable household contact (sibling/parent or adult caregiver with high risk condition or elderly) a delay in the return to school may be considered.
- Adults at schools (teachers, administrative and support staff) with underlying medical conditions
 or other risk factor for severe disease could potentially acquire COVID in the school or daycare
 setting. Additional precautions may be warranted and they should consult with their primary
 care provider and/or specialist.
- Prior to return to school for children with potential high risk condition consultation with their primary care physician or specialist may be warranted – to review clinical status and medications.
- Review of immunization status to avoid vaccine-preventable diseases.

In Summary

- There are tremendous benefits from children attending school.
- There are potential risks from children not attending school for prolonged periods of time.
- COVID-19 does not appear to cause severe disease in children
- Until broad molecular and antibody testing are available to better understand the seroprevalence of COVID-19 in our community, it may be prudent for a select group of patients to remain at home.
- The ultimate decision should be made by family, with support and guidance from their primary care team and/or subspecialist.

References:

- Lifting of restrictive public health measures Recommendations from the F/P/T Special Advisory Committee on COVID-19 (Accessed at: https://www.canada.ca/en/public-health-measures.html)
- 2. Update on COVID-19 epidemiology and impact on medical care in children: April 2020 (Accessed at: https://www.cps.ca/en/documents/position/update-on-covid-19-epidemiology-and-impact-on-medical-care-in-children-april-2020)
- 3. AAP guidance on school reopening addresses physical and mental health, instructional time (Accessed at:
 - $\frac{\text{https://www.aappublications.org/news/2020/05/covid19schoolopening050520)}{\text{https://services.aap.org/en/pages/2019-novel-coronavirus-covid-19-infections/covid-19-planning-considerations-return-to-in-person-education-in-schools/}$
- 4. Parri N, Lenge M, Buonsenso D; Coronavirus Infection in Pediatric Emergency Departments (CONFIDENCE) Research Group. Children with Covid-19 in Pediatric Emergency Departments in Italy. N Engl J Med. 2020 May 1.
- COVID-19 in schools—the experience in NSW. Prepared by the National Centre for Immunisation Research and Surveillance (NCIRS) 26 April 2020 (not peer-reviewed). (Accessed at: http://ncirs.org.au/sites/default/files/2020-04/NCIRS%20NSW%20Schools%20COVID_Summary_FINAL%20public_26%20April%202020.pdf



